



Co-funded by the Erasmus+ Programme of the European Union

ABOUT THE PROJECT

In September 2018, Jakub Krčín´s High School of Fisheries and Water Management (SŠRV) joined the Erasmus+ programme again. This has resulted in the three partner schools: SŠRV, Nord-Troms videregående skole and Lyceé Professional de Guérande - Olivier Guichard, once again joining together in the two-year international project Food from water Used To Urge Revolution in Eating habits (FUTURE). The school's project teams were tasked with concentrating their attention on aquaculture, the ecological aspects of sustainable healthy food production and the promotion of fish, seafood and freshwater products.

The first project meeting took place in November 2018 in Třeboň, where students concentrated on carp, grass carp and crayfish. In May 2019, the teams met in Skjervøy, Norway, to get a closer look at salmon, sea urchin and seaweed. The third student mobility took place in October 2019 in Guérande in France, where attention was drawn to oysters, green algae and to an introduction of the aquaponic system.

INVOLVED SCHOOLS



JAKUB KRČÍN'S HIGH SCHOOL OF FISHERIES AND WATER MANAGEMENT TŘEBOŇ CZECH REPUBLIC

Jakuba Krčín's High School of Fisheries and Water Management is located in Třeboň, a town close to the border in the south of the Czech Republic. Due to its untouched landscape and unique pond system, the region of Třeboň is a UNESCO World Heritage Site.

A professional fishing apprenticeship school was established in Třeboň in 1951, the only one in the entire Czechoslovakia. The fishing apprenticeship tradition was then followed by the Jakub Krčín´s High School of Fisheries and Water Management in 2008.



The Nord-Troms videregående skole – seek this school far beyond the Arctic Circle, around the 70th parallel, in the county of Tromso. The school was founded in 1990 in an area typical for farming Atlantic salmon, Alpine grayling and Rainbow trout. Today, it is not only involved in aquaculture and fishing, but also in fish gastronomy.



The Lyceé Professionel de Guérande is located near the Atlantic Ocean, in a tourist and agricultural area near Nantes. Founded in 1972, this school was the first in France to deal with water management. Today, it educates future experts in catering, landscape architecture or fish and shellfish farming.

eTwinning

You can find all of the project's outputs on: http://twinspace.etwinning.net/71124/home

The European Commission's support for the production of this publication does not constitute an endorsement of the contents, which reflect the views only of the authors, and the Commission cannot be held responsible for any use which may be made of the information contained therein.



SPECIES TYPICAL FOR THE CZECH REPUBLIC

CARP

(Cyprinus carpio)

HISTORY AND ORIGIN OF THE SPECIES

The common carp belongs to the evolutionary younger fish – the bony fish and according to recent knowledge carp appeared for the first time around the Black Sea and from there migrated and got to further places like Russia, China and the Czech Republic.

The first reference about carp breeding is around 1000BC in ancient China. Carp was considered as a symbol of luck and wealth and also was part of an important food source. It is also interesting that quite often there were colourful variations which we know nowadays as Koi carp. In Europe carp began spreading in the catchment area of the Danube River, nowadays we call this wild form 'Sazan'. The carp got into the catchment area of the Elbe River thanks to Romans who fished for it and used it as a food source for their soldiers. These soldiers were moving a lot so they needed to take carp with them. So the carp got into the Elbe River where it escaped the Romans and migrated further west. This was the way the carp got into the whole Europe. It was really popular around 1000AC when it was considered as a suitable fasting dish. Therefore the professional breeding was really started by the monasteries and then by aristocratic dynasties like the Pernštejns and Rosenbergs who helped spread carp further.

DESCRIPTION, BIOLOGY AND GROWTH

Carp is the most typical representative of Cyprinidae family all over the world. It has spindle-shaped form of the body with a distinctive spine. The height of the spine can differ according to genetic line or environment where it lives. The head is distinctive with pharyngeal jaws. A great recognizable sign is also a pair of whiskers/barbels/ at each corner of the mouth. The single dorsal fin is prolonged and simple. The forked tail fin is quite big with deep shape. The interesting part are the scales as thanks to genetic breeding we know four variations of scales on carp. They include large shiny scales, absence of scales, scales in a line along the flanks or smooth scales. Carp is a very easily satisfied species. It needs higher temperatures but it is also able to survive in a very cold waters.

Stagnant or running waters are suitable and this species is not liable to organic pressure. It even manages on a minimum concentration of oxygen. It creates a big shoal and is not aggressive to other fish even though it is actually omnivorous.

Growth is dependent on the amount of food and the water temperature. In the waters of the Czech Republic it grows up to weight of 2,5kg in about 3-4 years. But in the warm waters of Australia it would grow to this size in just over a year.

NATURAL HABITAT AND AOUACULTURE

Naturally the common carp only occurred in Asia. Today with human activity carp spread extremely and we can find it in almost any country.

It is a really flexible species which can prosper in rivers, dams, ponds and lakes or even in various small storage reservoirs. From these locations it also migrates to further distances. This way it is able to spread naturally practically anywhere.

Thanks to these qualities carp is either loved or hated. In USA waters or Australia carp is extremely overbred and is pushing other species out.

Opposite to that, in the Czech Republic it has been actively bred in ponds for hundreds of years. It is stocked in lakes, fed by corn and is much liked. It is the most bred species of fish and the Czech Republic is able to produce 17000t of carp a year which is the fourth largest production of carp in the world.

REPRODUCTION IN THE NATURE AND ARTIFICIAL REPRODUCTION

The carp in our conditions matures around 3-4 yrs of age. The male fish matures a year sooner. The mating usually takes place during the spring months when the water temperature is above 20°C. Both male and female carp don't migrate because of reproduction. They only look for parts close to the bank with plenty of vegetation. Carps put their fish eggs onto water plants. There has to be lots of plants because one female can produce 1.5 million fish eggs.

If all conditions are met, males start to follow females usually 2:1. One female would have at least two males and flow in a shape of an arrow. Then they all flow into the water plants where the females release the fish eggs and the male the sperm to fertilize the eggs. Mating is quite fierce but doesn't take a long time. Fertilized eggs stick to the plants where they stay for about three days.

Nowadays the reproduction of carp in nature happens very rarely because natural reproduction is not very effective and wouldn't cover the need of the breeders. So therefore today artificial reproduction is mainly used. That is worked out so well with the carp and it is not very difficult.

Mature and prepared fish are moved to incubators where they are weighed, sorted and marked. Twenty four hours before reproduction they are given hormones to help start reproduction. Most of time hypophysis – part of the carps brain is given by injection.

After 24 hours the fish are picked up and by massaging the belly we get fish eggs and sperm in clean bowls. After getting enough amount they are both mixed and covered with water. Then the fertilization starts and takes about 3-5 minutes. Fish eggs become very sticky which is undesirable so with the help of the milk fat it's possible to get rid of the stickiness.

Fertilized eggs are moved into special incubation bottles where they stay for 3 days before carp embryo is born.

QUALITY OF MEAT AND NUTRITIONAL TABLE

The description of carp meat isn't clear. Generally it is a dietetic meat, easy to digest and overall beneficial. But the quality is strongly influenced by the environment and type of breeding. It's possible to find a carp meat where quality and taste is perfect. But it is also possible to find a carp which is practically inedible.

In the Czech Republic carp has been bred for hundreds of years and its breeding is worked out in detail. That leads to a fact that the Czech carp has such a name as Norwegian salmon or French wine.

Sugars(g/100g)	7
Fats (g/100g)	10
Proteins (G/100g)	20
HUFA(%)	36
PUFA(%)	18
MUFA(%)	39
Omega 3 (%)	8
Vitamins	A,E,K
Energetic value (kJ/100g)	648

SIGNIFICANCE OF THE SPECIES NOWADAYS AND IN THE FUTURE

Carp is a very important species and belongs to the fifteen most bred freshwater fish in the world. Some breeding is entirely based on its production and wouldn't be possible to exist without it. It is also a truth that recently carp is not able to compete with other species especially seawater fish which are imported practically anywhere.

So it is presumed that the production of carp won't be growing in the future but will stay where it is now.



CRAYFISH

(Astacidea)



HISTORY AND ORIGIN OF THE SPECIES

There are about five species of crayfish in the Czech Republic. Recently there were also other species imported because of aquaristics and these species are now getting into our waters.

The crayfish were naturally part of our waters and their fishing and consumption were common There is a proof in past documents which mention that in 1910 there were over 500 official crayfish fisheries. The Natural habitat of crayfish is not limited only to nature parks or conservation areas where the quality of water is carefully looked after. Natural habitat is documented all over our country. Quite often it's invasive population of crayfish which is found as they are less demanding and push native crayfish out.

DESCRIPTION, BIOLOGY AND GROWTH

The body of the crayfish is formed by the head and joined thorax, abdomen and limbs. The top of the body is protected by chitinous carapace. Midsection is divided with diagonal nape line to two parts head and body, which became one part. Abdomen of the crayfish is formed by seven segments. The last of them turned into a tail fin / uropod. The abdomen is very flexible and mobile.

The crayfish have 19 pairs of limbs. That also include antennae, mandibles, claws, maxillipeds – out of which only five pairs of limbs are for walking.

Crayfish need a really good quality of water and lower temperatures. Organic presssure can cause problems and chemical strain is deadly. Often they are seen as indicators of the quality of water. Most crayfish are most active at night and spend most of the time being hidden. They are mainly herbivores but can eat bits of meat or even their own species.

Growth is atypical and dependent on molting of the shell. In optimal conditions crayfish can molt their shell 10-12 times in the first year of their life. Our heaviest crayfish can reach weights of 200g in 4-7 years.

NATURAL HABITAT AND AQUACULTURE

Natural habitat of crayfish all over the world is influenced by native or non native species. All crayfish need locations with very clean water, pebbly river bottoms, lots of hiding spaces, abundance of oxygen and a lot of food. These waters can usually be found only in mountain rivers in protected areas. Invasive crayfish have far less demands so can appear in less suitable waters where native crayfish wouldn't survive. So all over the world we can see the disappearance of native species of crayfish and the expansion of the others. The waters of the Czech Republic are no exception and the situation is getting worse due to importing crayfish for aquaristics, these crayfish then get out into the waters. For these reasons we create farms for crayfish which should function as rescue centres for native species, support of crayfish in their natural habitat and recently also forms for consumption. Crayfish

species, support of crayfish in their natural habitat and recently also forms for consumption. Crayfish live in earthy ponds or special tanks with an abundance of hiding spaces and are fed with special granules or carrots.

REPRODUCTION IN NATURE VERSUS ARTIFICIAL REPRODUCTION

Crayfish in the Czech Republic mate from September to October when they're 4 years old. During mating some mature males can migrate a bit. Reproduction happens through spermatophore, which is a bag of non active sperm which male sticks to a female. It's interesting that the female can decide whether to accept this bag or discard it. That's why one male can mate with 15 females.

About 2-3 weeks after mating the female starts to lay eggs. Depending on the species it could be between 40-550 eggs. During the laying, the spermatophore is also dissolved and activates the sperm. Then the eggs are fertilized. The female sticks the eggs under her abdomen and hides away to stay without any food. New crayfish are born in April.

During the first few days the newborns hide under the tail of their mother until the first or second molting of their shell. This happens within two weeks. After that the crayfish start to become independent and search for safety. They feed independently and carry on molting their shell.

Because of a bad situation there are lots of farms set up, which are using artificial reproduction to get a new generation of crayfish. It is interesting that even in these professional breeding places the crayfish have a very good way of looking after the descendants.

Artificial reproduction is very simple. A breeder just waits for the right time to put the crayfish together in special tanks and waits for the mating. Then he takes the females with their eggs and leaves them in the tank alone. Females look after the eggs really well, a breeder just need to check the quality of water.

The female tries to push away the hatched crayfish after four weeks as they are independent and the breeder has an ideal opportunity to take them away and move elsewhere.

QUALITY OF MEAT AND NUTRITIONAL TABLE

The quality of crayfish meat is very high and is compared with the quality of prawns, lobsters or crabs. This isn't by chance as all these mentioned above belong to crustaceans and therefore are very similar. The next proof of a high quality of crayfish meat could be in any historical documents, which talk about crayfish as an aristocratic food.

It is a dietetic meat, easily digestible and with no risk of disease transfers. The only exception could be a seafood allergy for some consumers which could potentionally happen after eating crayfish meat.

Last but not least it is also important to think about the fact that the original species of crayfish are protected and their consumption is a crime. Meat could be obtained only from non native species of crayfish

Sugars (g/100g)	3
Fats (g/100g)	2
Proteins (G/100g)	17
HUFA (%)	29
PUFA (%)	14
MUFA (%)	41
Omega 3 (%)	0
Vitamins	A,E,K
Energetic value (kJ/100g)	591

SIGNIFICANCE OF THE SPECIES NOWADAYS AND IN THE FUTURE

Nowadays the crayfish is considered to be an endangered species so it is really necessary to keep an eye on their protection and survival. Unfortunately the recent studies show that despite all the efforts the original species of crayfish is pushed out by non native species all the time. In many countries there is no need to save the native species as the non native species has taken over the waters.

This way the population of non native species is still rising and become more available. In the USA or in France it is common to eat crayfish in restaurants or homes. And also crayfish meat becomes more popular all the time even though it will always be just non native species..



HISTORY AND ORIGIN OF THE SPECIES

Amur is native to Asia but was introduced to many parts of the world including Europe and the Czech Republic. It also appears in Africa and the USA. Originally amur was only found around the Amur River in the east of Russia and naturally migrated to China.

It was mainly a river species living in the big lowland rivers and their branches. It was imported into our country after 1960 from the Soviet Union, mainly to Třeboň and its job was to naturally eliminate higher vegetation in water. Also because of the political situation of the world in the sixties we could debate about the political reasons of bringing amur to the Czech Republic.

DESCRIPTION, BIOLOGY AND GROWTH

It's a typical fish of the Cyprinidae family which is quite often confused with carp by the public. But there are some typical differences to a carp. Amur hasn't got any whiskers on both sides of the mouth. Its head is wedge-shaped and flat on the top. The position of the eyes is low and there is a distinctive round dorsal fin. But the golden colour of its sides and white belly is the same as carp.

Amurs live in big shoals in warm and deep waters and feed on higher plants. They are called 'living lawnmowers' as they manage to eat abnormal amounts of high plants. Sometimes it's like 50kg of reed to 1kg of weight.

The sizes can sometimes appear abnormal. Some resources talk about weights of 30-45kg. This extreme growth could only show with older fish which can add every year 2kg. Young fish grow slowly and a year old amur's weight is only 5g.

NATURAL HABITAT AND AQUACULTURE

The natural home of this species is the Amur River which is also classed as one of the biggest rivers in the world which also separates countries like China and Russia. Thanks to humans we can find amurs in many parts of the world. The exceptions aren't states like USA, Canada, JAR, Hungary, Austria and the Czech Republic. The natural habitat is divided into two camps. In some countries the species is accepted as positive and in other countries it could be classed as an extremely invasive species which is pushing other species out. A scary example is the over reproduction of Amur in North America where they were stocked as a weed control in canals. The situation is out of control and massive shoals of amurs are pushing other species out. But even with using electricity or special poisons for fish, they still can't get this under their control.

Opposite to that a good example is breeding amurs with carps which is done in Czech ponds. Carps and amurs need similar environment so they can be bred together. Also worldwide amur is quite a favourite so it belongs to five most bred freshwater fish.

REPRODUCTION IN NATURE AND ARTIFICIAL REPRODUCTION

Reproduction of this species is totally connected with the Amur River which gives fish ideal conditions for reproduction. The fish ready for reproduction create massive shoals which are 10-15m long. Females start to release fish eggs. It's interesting that these eggs contain a big amount of fat which helps their floating. Also every female can release up to hundreds of thousands of fish eggs. Then the males flow through a cloud of eggs and release the sperm which fertilize the eggs.

Amur is really a warm water lover and in many rivers where they was imported, they can't get these temperatures. It needs temperatures over 25°C. This affects its reproduction which is then very limited. So it's a fish which can't naturally reproduce in the Czech Republic.

So the Czech Republic is 100% dependent on artificial reproduction. But the good news is that this species has the easiest artificial reproduction out of all the fish in our waters. It's important to follow up several basic information and procedures.

Amurs are mature around 5-7 years old. The males mature sooner. The water temperature shouldn't fall under 23°C. So the artificial reproduction is done between the end of June and beginning of July. The artificial reproduction starts with dividing the male and female fish and giving them hormones. Most of time it is hypophysis given by injection which comes from carp brains and is given 24hours before the planned mating. The fish is kept separated so it doesn't go into spontaneous reproduction.

After 24hours the fish is picked up and by massaging its belly we get fish eggs and sperm into clean bowls. After getting enough of the amount it is all mixed together and water is poured over it. Development of embryo lasts 30hours and is finished by leaving the fish egg.

QUALITY OF MEAT AND NUTRITIONAL TABLE

In recent years there are a lot of people in the fishing industry who are in agreement about the high quality of amur's meat. Even in sensory tests amur's winning against other fish of Cyprinidae family, mainly carp itself. The consumption of amur's meat grows every year and some study say that one day amur could compete with carp even during Christmas sale.

Sugars (g/100g)	5
Fats (g/100g)	4
Proteins (G/100g)	18
HUFA (%)	33
PUFA (%)	18
MUFA (%)	45
Omega 3 (%)	4
Vitamins	A,E,K
Energetic value (kJ/100g)	638

SIGNIFICANCE OF THE SPECIES NOWADAYS AND IN THE FUTURE

The significance of amur is today divided between positive and negative. In areas, where it's popular, the popularity grows every year. Because of its meat and for sports fishing we expect that popularity will grow in the future and this species will belong to three mainly bred freshwater fish in the world. In opposite areas where amurs are causing problems it is expected that even with all the effort it won't be possible to manage the situation. The problems will grow and lead to elimination of other species of fish.

In the Czech Republic amur is a real favourite and it would be good to slowly raise its production.



HEALTHY FOOD - SELECTED SPECIES

SPECIES TYPICAL FOR FRANCE


OYSTER


(Ostrea edulis)

HISTORY AND ORIGIN OF THE SPECIES

Oysters has always been eaten since Ancient times. The culture of the indigenous flat oyster (Ostrea edulis) began in France in the 17th century. The oyster beds were overexploited from the 18th century, especially along the Atlantic coast and the landings became poor and irregular during the 19th century. So, in 1860, the Portuguese cupped oyster (Crassostrea angulata) was chosen to replace the flat oyster until the 1970s, when a disease led to the total extinction of the Crassostrea angulata in France. It was then replaced by the Japanese cupped oyster (Crassostrea gigas) which was more resistant to the disease. Nowadays it is the most cultivated species in France and worldwide.

DESCRIPTION, BIOLOGY AND GROWTH





Oysters are bivalve molluscs which live in salt water. The optimal temperature for the growth is 21°C. Their diet is based on phytoplankton. Oysters are filter feeders, drawing water in over their gills through the beating of cilia. Suspended plankton and particles are trapped in the mucus of a gill, and from there are transported to the mouth, where they are eaten.

NATURAL HABITAT AND AQUACULTURE


Different production techniques according to the rearing area:

Along the Atlantic coast: Off-bottom culture is done by using plastic mesh bags set on trestles (tables) in the intertidal zone.

Along the Mediterranean coast: Suspended culture (5%), is done by hanging oysters fixed on ropes or in baskets from special frames (tables).

REPRODUCTION IN NATURE VERSUS ARTIFICIAL REPRODUCTION

approximately 2 weeks



QUALITY OF MEAT AND NUTRITIONAL TABLE







Amount Per 100 gran	ms ▼		
Calories 163			
			% Daily Value
Total Fat 4.6 g			7%
Saturated fat 1	g		5%
Polyunsaturated	i fat 1.8 g		
Monounsaturate	d fat 0.8 g		
Cholesterol 100 mg			33%
Sodium 212 mg			8%
Potassium 302 mg			8%
Total Carbohydrate	10 g		3%
Dietary fiber 0 g			0%
Sugar 0 g			
Protein 19 g			38%
Vitamin A	9%	Vitamin C	21%
Calcium	1%	Iron	51%
Vitamin D	0%	Vitamin B-6	5%
Vitamin B-12	479%	Magnesium	11%

SIGNIFICANCE OF THE SPECIES NOWADAYS AND IN THE FUTURE

We eat oysters more particularly during Chistmas and New Year period. It is a tradition in most French families. They are really healthy since they are rich in omega 3 fatty acids and trace elements important for the body (skeletton, blood). Once you have understood how to open them, it is a real pleasure to share a plate of oysters with your friends!